

Brighton[®]



AGT

Variety snapshot

- Dual purpose winter wheat, suitable for grazing and grain production
- A higher yielding alternative to Illabo^Φ, EGA Wedgetail^Φ and LRPB Kittyhawk^Φ
- Mid winter maturity, similar to Illabo^Φ
- Improved test weight compared with Illabo^Φ and EGA Wedgetail^Φ
- Improved yellow leaf spot resistance over Illabo^Φ and EGA Wedgetail^Φ
- AH quality classification in Vic/SA

Breeder's comments

Dual purpose, graze and grain wheat varieties have traditionally been very valuable to mixed farmers, providing more than one opportunity to generate income throughout the season. The use of dual purpose varieties has continued to gain in popularity, perhaps due to a shift in earlier sowing of grain-only crops.

We started a winter wheat breeding program at Wagga Wagga in 2014 in acknowledgement of the need for better performing long season and dual purpose varieties, with Illabo[®] being a popular release from this program. Illabo[®] has been a success story, offering mixed farmers a large step up in performance over the mainstay variety EGA Wedgetail[®].

Our newest variety in this space, Brighton[®], is poised to offer even more advancements in productivity, offering improvements in yield and physical grain quality over Illabo[®].

Brighton[®] also offers improved yellow leaf spot resistance over Illabo[®], however is more susceptible to powdery mildew and septoria tritici blotch.

Brighton[®] is a mid maturing winter wheat, reaching head emergence at a similar time to Illabo[®] across a range of sowing dates.

Brighton[®] is derived from popular main season wheat variety Beckom[®], and has inherited Beckom's[®] shorter plant height and aluminium (acid soils) tolerance genes. Like Beckom[®] and many other varieties, Brighton[®] may express physiological leaf yellowing throughout winter; however will grow out of these symptoms in spring.

To maximise grain only yield, Brighton[®] appears ideally suited to mid-late April sowing in high yield environments, and mid April in lower yielding environments. To maximise the length of safe grazing time, Brighton[®] may be sown from mid March through to mid April.

Table 1. Specifications

Background

Tested as	V14051-172
Released	2024
EPR rate	\$4.10/tonne + GST

Disease

Stem Rust resistance*	MRMS
Stripe Rust resistance*	MRMS
Leaf Rust resistance*	S
Yellow Leaf Spot resistance*	MRMS
Powdery Mildew resistance* (SA)	SVS
Powdery Mildew resistance* (VIC)	MSS (P)
Septoria Tritici Blotch resistance*	S
CCN resistance*	R
Pratylenchus Neglectus resistance*	S
Pratylenchus Neglectus tolerance*	IVI
Eyespot resistance*	S
Crown Rot resistance*	S

Plant Characteristics

Maturity speed^	Mid
Maturity habit^	Winter
Sowing window^	Early
Novel herbicide tolerance^	None (conventional tolerance)
Head type^	Awned
Plant height^	Short-moderately short
Coleoptile length^	Moderate
Lodging tolerance^	MT

Abiotic Stress

Boron tolerance^	Does not carry tolerance gene
Acid/aluminium tolerance^	Carries tolerance gene

Grain Quality

Quality classification	AH
Grain colour	White
Screenings level^	Very low
Test weight^	Very high
Sprouting tolerance^o	I
Black Point resistance*	MS

Legend


R	Resistant	VI	Very Intolerant	o	Rating based on Germination Index Values
MR	Moderately Resistant	(P)	Provisional rating	^	AGT ratings/data interpretation. Comprehensive AGT agronomic trait ratings and data can be found at: https://bit.ly/Traits_2026
MS	Moderately Susceptible	NA	Not Available		
S	Susceptible	/	Pathotype differences		
VS	Very Susceptible	-	Range		
T	Tolerant	,	Mixed phenotype		
MT	Moderately Tolerant	#	May be more susceptible to alternate pathotypes		
MI	Moderately Intolerant	*	NVT consensus ratings 2026		
I	Intolerant				

Grain yield

In both long season NVT trials (Figure 1) and early sown NVT trials (Figure 2) across higher rainfall zones of SA/Vic, Brighton[®] has produced slightly higher yields than Illabo[®] over-all, and much higher yields than EGA Wedgetail[®] and LRPB Kittyhawk[®].

NVT long season trials are generally sown in April, in areas that experience softer, elongated springs. NVT early sown trials are generally sown in late April/early May.

Figure 1. Grain yield of Brighton[®] versus comparators across SA/VIC - NVT long season trials



Source: Best linear unbiased predictions from NVT long term MET analysis, long season trial series 2021-2025


[] : Total number of trials per region

() : Number of trials that each variety was present in across the dataset

{ } : Region mean

Grain yield

Figure 2. Grain yield of Brighton^ϕ versus comparators across SA/VIC - NVT early sown trials



Source: Best linear unbiased predictions from NVT long term MET analysis, early sown trial series 2021-2025
[]: Total number of trials per region
(): Number of trials that each variety was present in across the dataset
{ }: Region mean

Variety comparisons

Brighton^ϕ has an AH quality classification in SA/Vic, and offers low screenings levels and a test weight improvement over main competitor Illabo^ϕ.

Brighton^ϕ offers a good disease resistance package, including good stripe rust resistance and an improvement in yellow leaf spot resistance compared with Illabo^ϕ.

Brighton's^ϕ powdery mildew and septorial tritici blotch resistance is lower than Illabo^ϕ.

Table 2. Variety comparisons

		Brighton ^ϕ	EGA Wedgetail ^ϕ	Illabo ^ϕ	LRPB Kittyhawk ^ϕ
Disease	Stem Rust resistance*	MRMS	MRMS	MR	MRMS
	Stripe Rust resistance*	MRMS	MS	MRMS	MR
	Leaf Rust resistance*	S	MS	S	MR
	Yellow Leaf Spot resistance*	MRMS	MSS	MS	MRMS
	Powdery Mildew resistance* (SA)	SVS	S	RMR	MS
	Powdery Mildew resistance* (VIC)	MSS (P)	MS (P)	MR (P)	MSS (P)
	Septoria Tritici Blotch resistance*	S	MSS	MSS	MRMS
	CCN resistance*	R	S	MRMS	S
	Pratylenchus Neglectus resistance*	S	S	MSS	S
	Pratylenchus Neglectus tolerance*	IVI	MII	MI	MI
	Eyespot resistance*	S	NA	S	S
	Crown Rot resistance*	S	S	S	SVS
Plant Characteristics	Maturity speed [^]	Mid	Mid	Mid	Mid
	Maturity habit [^]	Winter	Winter	Winter	Winter
	Sowing window [^]	Early	Early	Early	Early
	Novel herbicide tolerance [^]	None (conventional tolerance)	None (conventional tolerance)	None (conventional tolerance)	None (conventional tolerance)
	Head type [^]	Awned	Awned	Awned	Awned
	Plant height [^]	Short-moderately short	Moderately short	Short-moderately short	Moderately short
	Coleoptile length [^]	Moderate	Short	Short	Short
	Lodging tolerance [^]	MT	MI	MT	MTMI
Abiotic Stress	Boron tolerance [^]	Does not carry tolerance gene	Does not carry tolerance gene	Does not carry tolerance gene	Does not carry tolerance gene
	Acid/aluminium tolerance [^]	Carries tolerance gene	Carries tolerance gene	Does not carry tolerance gene	Does not carry tolerance gene
Grain Quality	Quality classification	AH	APW	AH	AH
	Grain colour	White	White	White	White
	Screenings level [^]	Very low	Low	Very low	Low
	Test weight [^]	Very high	Very low	Low	Very high
	Sprouting tolerance ^{^ϕ}	I	MII	MII	I
	Black Point resistance*	MS	MS	MRMS	MRMS



Seed Availability

Please contact an AGT Affiliate or your local retailer for seed. Consult the AGT website for AGT Affiliate contact details (www.agtbreeding.com.au/affiliates). AGT varieties can be traded between growers upon the completion of a License Agreement as part of AGT's Seed Sharing™ initiative (www.agtbreeding.com.au/seedsharing)

PBR and EPR

Varieties denoted by the [®] symbol are protected by Plant Breeders Rights (PBR) and all production (except seed saved for planting) is liable to an End Point Royalty (EPR), which funds future plant breeding. Growers of PBR protected varieties will be subject to a Grower License Agreement that acknowledges that an EPR must be paid on all production other than seed saved for planting.

Contact

Brad Koster, Variety Support Manager, SA:	0400 812 475
Rowan Prior, Variety Support Manager, Vic:	0429 677 736
AGT End Point Royalty team:	(08) 7111 0201
agtbreeding.com.au	

The information contained in this brochure is based on knowledge and understanding at the time of writing. Growers should be aware of the need to regularly consult with their advisors on local conditions and currency of information. Wherever possible, independent NVT data has been used in this publication. In the absense of NVT data, AGT data has been provided.